

are other SOx proteins responsible for S-S that remain undetected by our antibody. We need further studies to obtain the c-DNA clone of SOx to detect the other isoforms of the enzyme in the keratinocyte cDNA library.

Yukiko Hashimoto, Yasushi Suga, Shouichi Matsuba,  
Masayuki Mizoguchi, Kenji Takamori, Juergen Seitz,\*  
Hideoki Ogawa

Department of Dermatology,  
Juntendo University School of Medicine, Hongo,  
Bunkyo-ku, Tokyo, Japan

\*Department of Anatomy and Cell Biology,  
Philipps University of Marburg,  
Marburg, Germany

## REFERENCES

- Danno K, Horio T: Histochemical staining of sunburn cells for sulphhydryl and disulphide groups: a time course study. *Br J Dermatol* 102:535–539, 1980
- Darmon M, Blumenberg M: Retinoic acid in epithelial and epidermal differentiation. In: Darmon M, Blumenberg M, eds. *Molecular Biology of the Skin*. San Diego: Academic Press, 1993; pp 181–206
- Goldsmith LA: The epidermal cell periphery. In: Goldsmith LA, ed. *Biochemistry and Physiology of the Skin*. New York: Oxford University Press, 1983; pp 197–199
- Hashimoto Y, Suga Y, Matsuba S, Mizoguchi M, Takamori K, Seitz J, Ogawa H: Immunofluorescent localization of sulphhydryl oxidase correlates with disulfide cross-linking in the upper epidermis of rat skin. *Arch Dermatol Res* 292:570–572, 2000
- Haugaard N: Cellular mechanisms of oxygen toxicity. *Physiol Rev* 48:311–373, 1968
- Kumari M, Aumüller G, Bergmann M, Meinhardt A, Seitz J: Stage-dependent appearance of sulphhydryl oxidase during spermatogenesis in the testis of rat and hamster. An immunohistochemical study. *Histochemistry* 94:365–371, 1990
- Ogawa H, Taneda A, Kanaoka Y, Sekine T: The histochemical distribution of protein bound sulphhydryl groups in human epidermis by the new staining method. *J Histochem Cytochem* 27:942–946, 1979
- Takahashi H, Iizuka H: Nucleotide analysis of the 5'-upstream promoter region of human involucrin gene: activation by 12-O-tetradecanoylphorbol-13-acetate. *J Invest Dermatol* 100:10–15, 1993
- Takamori K, Thorpe JM, Goldsmith LA: Skin sulphhydryl oxidase. Purification and some properties. *Biochem Biophys Acta* 615:309–323, 1980
- Yamada H, Takamori K, Ogawa H: Localization and some properties of skin sulphhydryl oxidase. *Arch Dermatol Res* 279:194–197, 1987
- Yamamoto K, Takamitsu S, Kanaoka Y: Fluorescent thiol reagents. XII. Fluorescent tracer method for protein SH groups using N-(7-dimethylamino-4-methyl coumarinyl) maleimide. An application to the proteins separated by SDS-polyacrylamide gel electrophoresis. *Anal Biochem* 79:83–94, 1977

# Abundant Expression of Vasoactive Intestinal Polypeptide Receptor VPAC<sub>2</sub> mRNA in Human Skin

To the Editor:

Vasoactive intestinal polypeptide (VIP) is a bioactive peptide that influences many aspects of cell function and differentiation. The 28-amino acid polypeptide belongs to the glucagon/secretin superfamily and is widely expressed in the central nervous system and in peripheral tissues including lung and skin, where it has been shown to have a multitude of biological functions (Dickinson and Fleetwood-Walker, 1999). VIP is abundantly present in cutaneous autonomic and sensory nerve fibers (Eedy *et al*, 1994) where it acts as neuromodulator and participates in the regulation of regional blood flow. As the expression and localization of VPAC<sub>2</sub> mRNA has not been examined so far, this study was carried out to correlate VIP and its receptor in the human skin. Surgically resected human skin samples (n = 24) were obtained from surplus remnants in excess of that required for pathologic examination after informed consent.

For mRNA *in situ* hybridization, VPAC<sub>2</sub> receptor cRNA probes were generated by a standardized protocol using the sequence corresponding to the region spanning from TM 3–7 of the human VPAC<sub>2</sub> receptor from TSUP1 human T lymphoblast cDNA which was subcloned into a specific vector (pGEM-T vector, Promega, Madison, WI) (Groneberg *et al*, 2001a). For antisense probes, plasmid linearization by restriction with Spe I was followed by transcription with T7 polymerase, for sense probes, the Nco I linearized plasmids were transcribed with SP6 polymerase (all Roche Diagnostics, Mannheim, Germany). The probes were checked for their integrity by TAE-agarose gel electrophoresis and ethidium bromide staining.

The VPAC<sub>2</sub> receptor mRNA distribution was assessed by nonisotopic *in situ* hybridization using a standardized protocol

(Groneberg *et al*, 2001b) with 6–8 µm cryostat sections of 4% paraformaldehyde fixed biopsies.

High resolution interference contrast microscopy revealed abundant staining in all areas of the biopsies (**Fig 1**). Positive VPAC<sub>2</sub> mRNA-signals were localized to the cytoplasm of keratinocytes with a signal intensity that was maximal at the basal zone and decreasing to superficial layers (**Fig 1A**). In the deep part of the dermis, VPAC<sub>2</sub> mRNA hybridization signals were present in cells of eccrine sweat glands and in cells of the germinative epithelium, matrix, and medulla of the hair follicle (**Fig 1C**). Also, positive VPAC<sub>2</sub> mRNA hybridization signals were localized to endothelial and mononuclear immune cells. Control hybridizations on alternate sections with equivalent amounts of the digoxigenin labeled sense-probe using the same hybridization conditions and washing stringencies were unstained (**Fig 1B, D**).


To relate the distribution of VPAC<sub>2</sub> mRNA to its ligand VIP, immunohistochemical studies were carried out using mouse polyclonal (1:1000, Biogenesis, Poole, U.K.) or monoclonal (1:100 Charles River, Southbridge, MA) VIP-antibodies as described before (Fischer *et al*, 1998) with 6–8 µm cryostat sections and antimouse fluorescein-5-isocyanate antiserum (1:400, Amersham, Braunschweig, Germany) and anti-rabbit biotin (1:200) – streptavidin Texas Red (1:400, Amersham) as secondary antibodies.

Fluorescence microscopy demonstrated an abundant expression of VIP-immunoreactivity in nerve fibers which were found as branching networks surrounding blood vessels, eccrine sweat glands, and hair follicles (**Fig 1E and F**) and in direct contact to VPAC<sub>2</sub> mRNA positive cells. Controls by omitting the primary or secondary antibodies and incubation with the preimmune serum did not reveal specific immunosignals.

VIP binding sites have so far been shown in sweat glands using unspecific binding techniques (Heinz-Erian *et al*, 1986) and in allergic contact dermatitis (Lundeberg and Nordlind, 1999). There is also a large body of evidence on the functional role of VIP as a possible sensory neuropeptide in normal skin and dermatologic disorders. VIP protein expression has been demonstrated in normal skin of young (Eedy *et al*, 1994) and elderly (Abdel-Rahman *et al*,

**Figure 1. Localization of VPAC<sub>2</sub> receptor messenger RNA and VIP immunoreactivity.**

Eight micrometer cryostat sections of human skin were subjected to nonradioactive *in situ* hybridization for VPAC<sub>2</sub> mRNA. VPAC<sub>2</sub> mRNA hybridization signals are localized to epidermal keratinocytes (A) and cells of the germinative epithelium, matrix, and medulla of the hair follicle (C). Control slides, treated with the sense probe, did not show specific staining (B, D). Immunohistochemistry for VIP resulted in abundant staining of VIP-immunoreactive nerve fibers that were present around sweat glands (E) and in the subcutis (F). Scale bars: (A, B) 160  $\mu$ m, (C, D) 60  $\mu$ m, (E, F) 40  $\mu$ m.



1992) people and in dermatologic disorders such as nodular prurigo (Abadia Molina *et al*, 1992) or allergic contact dermatitis (Lundeberg and Nordlind, 1999). Also the functional role of VIP in the normal skin and in disorders like contact dermatitis (Bondesson *et al*, 1996) was analyzed; however, a detailed study of the mRNA expression of molecular distinct VIP receptors has not been carried out.

As the molecular properties of the VIP receptors were identified in the past years, this study was designed to localize the mRNA of the inducible VIP receptor VPAC<sub>2</sub> in normal human skin biopsies and resulted in abundant staining for VPAC<sub>2</sub> mRNA in keratinocytes. Effects of VIP on keratinocyte cultures have been demonstrated earlier *in vitro* (Haegerstrand *et al*, 1989). This demonstration of VPAC<sub>2</sub> receptor mRNA in human keratinocytes *in situ* supports the previous *in vitro* findings and provides evidence that the mode of VIP modulation of cell proliferation in keratinocytes is mediated at least partly via the VPAC<sub>2</sub> receptor. This finding is also supported by the recent demonstration of VPAC<sub>2</sub> immunoreactivity in allergic contact dermatitis (Lundeberg and Nordlind, 1999).

A dense network of VIP-IR nerve fibers around sweat glands and hair follicles was shown in this study, and VIP has been reported to induce cAMP generation in human sweat glands (Tainio, 1987). By directly localizing VPAC<sub>2</sub> mRNA to eccrine gland and hair follicle cells we provide first evidence for the identity of the VIP receptor subtype that is involved in VIP-induced upregulation of eccrine sweat secretion.

VIP has been shown to be a potent relaxant of vascular smooth muscle (Lundberg *et al*, 1981). As VPAC<sub>2</sub> receptor mRNA was not present in the vascular smooth muscle layer of cutaneous vessels, our results indicate that VIP-induced dermal vasodilation is mediated via a different VIP receptor or by paracrine VPAC<sub>2</sub> stimulation. Also, the demonstration of VPAC<sub>2</sub> mRNA in mononuclear inflammatory cells is a novel finding with significant

impact on the role of VIP in cutaneous immunomodulation, as a multitude of functional studies have suggested an important role of the peptide in the regulation of the immune system (Bellinger *et al*, 1996).

In conclusion, this study demonstrates an abundant expression of VPAC<sub>2</sub> mRNA in human skin that is associated to VIP immunoreactive nerve fibers and suggests a major participation of the VPAC<sub>2</sub> receptor in cutaneous VIP-signaling.

*We thank E. J. Goetzl (San Francisco) for generously providing the plasmids. This study was supported by the DFG (SFB 549, C1).*

Tanja C. Fischer,\*† Petra Hartmann,\* Christoph Löser,‡  
Jochen Springer,\*§ Christian Peiser,§ Q. Thai Dinh,\*§

Axel Fischer,§ David A. Groneberg\*§

\*Institute of Anatomy and Cell Biology,

Justus-Liebig-University, Giessen, Germany

†Department of Dermatology, Charité, Humboldt-University,  
Berlin, Germany

‡Department of Dermatology, Faculty of Medicine,  
Justus-Liebig-University, Giessen, Germany

§Division of Allergy Research,

Department of Pediatric Pneumology and Immunology, Charité,  
Humboldt-University, Berlin, Germany

## REFERENCES

- Abadia Molina F, Burrows NP, Jones RR, Terenghi G, Polak JM: Increased sensory neuropeptides in nodular prurigo: a quantitative immunohistochemical analysis. *Br J Dermatol* 127:344-351, 1992
- Abdel-Rahman TA, Collins KJ, Cowen T, Rustin M: Immunohistochemical,

- morphological and functional changes in the peripheral sudomotor neuro-effector system in elderly people. *J Auton Nerv Syst* 37:187-197, 1992
- Bellinger DL, Lorton D, Brouxhon S, Felten S, Felten DL: The significance of vasoactive intestinal polypeptide (VIP) in immunomodulation. *Adv Neuroimmunol* 6:5-27, 1996
- Bondesson L, Nordlind K, Mutt V, Liden S: Vasoactive intestinal polypeptide inhibits the established allergic contact dermatitis in humans. *Ann N Y Acad Sci* 805:702-707, 1996
- Dickinson T, Fleetwood-Walker SM: VIP & PACAP: very important in pain. *Trends Pharmacol Sci* 20:324-329, 1999
- Eedy DJ, Shaw C, Johnston CF, Buchanan KD: The regional distribution of neuropeptides in human skin as assessed by radioimmunoassay and high-performance liquid chromatography. *Clin Exp Dermatol* 19:463-472, 1994
- Fischer A, Canning BJ, Udem BJ, Kummer W: Evidence for an esophageal origin of VIP-IR and NO synthase-IR nerves innervating the guinea pig trachealis: a retrograde neuronal tracing and immunohistochemical analysis. *J Comp Neurol* 394:326-334, 1998
- Groneberg DA, Hartmann P, Dinh QT, Fischer A: Expression and distribution of vasoactive intestinal polypeptide receptor VPAC2 mRNA in human airways. *Lab Invest* 81:749-755, 2001a
- Groneberg DA, Nickolaus M, Springer J *et al*: Localization of the peptide transporter PEPT2 in the Lung: implications for pulmonary oligopeptide uptake. *Am J Pathol* 158:707-714, 2001b
- Haegerstrand A, Jonzon B, Dalsgaard CJ, Nilsson J: Vasoactive intestinal polypeptide stimulates cell proliferation and adenylate cyclase activity of cultured human keratinocytes. *Proc Natl Acad Sci USA* 86:5993-5996, 1989
- Heinz-Erian P, Paul S, Said SI: Receptors for vasoactive intestinal peptide on isolated human sweat glands. *General Pharmacol* 17:321-326, 1986
- Lundberg JM, Anggard A, Emson P, Fahrenkrug J, Hokfelt T: Vasoactive intestinal polypeptide and cholinergic mechanisms in cat nasal mucosa: studies on choline acetyltransferase and release of vasoactive intestinal polypeptide. *Proc Natl Acad Sci USA* 78:5255-5259, 1981
- Lundeberg L, Nordlind K: Vasoactive intestinal polypeptide in allergic contact dermatitis. an immunohistochemical and radioimmunoassay study. *Arch Dermatol Res* 291:201-206, 1999
- Tainio H: Cytochemical localization of VIP-stimulated adenylate cyclase activity in human sweat glands. *Dev Biol* 123:179-190, 1987